

## Differential Diagnosis of Microcytic Anemia, Thalassemia or Iron Deficiency Anemia: A Diagnostic Test Accuracy Meta-Analysis

Mina Jahangiri<sup>1</sup>, Fakhre Rahim<sup>2,3</sup>, Amal Saki Malehi<sup>1</sup>  
Seyed Mohammad Sadegh Pezeshki<sup>2</sup>, Mina Ebrahimi<sup>2</sup>

1. Department of Biostatistics and Epidemiology, Faculty of Health, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

2. Thalassemia and Hemoglobinopathy Research Center, Research Institute of Health, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

3. Metabolomics and Genomics Research Center, Endocrinology and Metabolism Molecular-Cellular Sciences Institute, Tehran University of Medical Sciences, Tehran, Iran

### KEYWORDS

Iron deficiency anemia;  
Thalassemia;  
Systematic review;  
Meta-analysis;  
Diagnostic test accuracy  
(DTA)

### Article Info

Received 2019/02/22;  
Accepted 2019/03/18;  
Published Online 2020

### ABSTRACT

We evaluated the most common indices to compare their sensitivity and specificity to introduce the most sensitive and specific index. We systematically searched five international indexing databases up to Dec 2018. For each index, we measured the diagnostic odds ratio (DOR), as well as summary ROC (SROC) curve which was used to compare the performance of each index. Deeks' tests of all discriminant indices indicated that there is no potential publication bias. The area under curves (AUCs) of all discriminant indices indicate overall good differential performance. The M/H ratio index was more sensitive and specific compared to other studied indices. In this meta-analysis, the M/H ratio index was more potential to discriminate iron deficiency anemia (IDA) from thalassemia trait. However, we cannot use this index alone to achieve the final diagnosis. The capability of this index to discriminate IDA from thalassemia trait must be used alongside with the common laboratory procedure to ensure the final differentiation.

**Corresponding Information:** Fakhre Rahim., Thalassemia and Hemoglobinopathy Research Center, Research Institute of Health, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran, E-mail: [bioinfo2003@gmail.com](mailto:bioinfo2003@gmail.com)

Copyright © 2020. This is an open-access article distributed under the terms of the Creative Commons Attribution-noncommercial 4.0 International License which permits copy and redistribute the material just in noncommercial usages, provided the original work is properly cited.

### Abbreviations

IDA, Iron deficiency anemia; ISI, Information Sciences Institute; RBCs, red blood cells; CBC, complete blood count; TP, True Positive; FP, False Positive; FN, False Negative; TN, True Negative; PLR, Positive Likelihood Ratio; NLR, Negative Likelihood Ratio; DOR, Diagnostic Odds Ratio; AUC, Area Under Curve; SROC, Summary Receiver Operating Characteristic.

### Introduction

Anemia is a common hematologic disorder and its lack of attention is associated with several physical and neurological complications [1]. On the other hand, its diagnosis and treatment are not very costly. The most common type of anemia is the microcytic anemia, of which two most common forms include Iron deficiency anemia (IDA) and  $\beta$ -thalassemia minor [2]. IDA is a common complication all across the world. Despite its high prevalence, there is still

some uncertainty in diagnosis steps [3]. Thalassemia is the most common monogenic disorder which is highly prevalent in the Mediterranean, Middle East, Indian peninsula and southeast of Asia. Nowadays, in the wake of extensive immigration in the past decades, thalassemia can be seen in almost everywhere [4]. Both IDA and thalassemia trait are among microcytic hypochromic anemias. In this type of anemia, globin chain synthesis disorders and heme

## 17 Diagnosis of Microcytic Anemia, Thalassemia or Iron Deficiency Anemia....

synthesis disorders cause microcytic and hypochromic red blood cells (RBCs), respectively [5]. Discriminating IDA and thalassemia trait in patients is a major challenge due to the resemblance of the anemic view. Evaluation of different hemoglobin levels by hemoglobin electrophoresis alongside with routine complete blood count (CBC) test and iron panel tests (such as ferritin level) are currently the main tools to discriminate IDA from thalassemia trait [6]. Despite their great utility, all routine hematologic tests such as HbA2, examination of a peripheral blood film, serum ferritin, iron, TIBC, and transferrin saturation are either not available in all clinical setups, or these are relatively time-consuming and expensive techniques. Consequently,

using these indices will save time and reduce diagnostic expenses [7, 8].

Due to the importance of discriminating these two complications in patient management and considering the financial limitations specifically in countries with high prevalence of thalassemia, mathematical indices, which are simpler and less complicated solutions, have been used to achieve a differential diagnosis. Despite the introduction of a considerable number of these indices in the past years, none of them are fully sensitive and specific to discriminate IDA from thalassemia trait. In this meta-analysis, we will evaluate the most common indices to compare their sensitivity and specificity in order to introduce the most sensitive and specific index.

**Table 1.** Discriminant indices for differentiation between thalassemia trait and iron deficiency anemia with their cut off values and references of publications based on these discriminant indices.

Discriminant Index	Formula	Cut off value		Number of studies	Number of patients	Ref.
		TT	IDA			
Ehsani [9]	$MCV - 10 \times RBC$	< 15	> 15	21	10817	[10-30]
England & Fraser (E&F) [31]	$MCV - RBC - 5 \times HB - 3.4$	< 0	0 >	81	26091	[10, 12-29, 32-90]
Green & King (G&K) [91]	$(MCV^2 \times RDW) / (100 \times HB)$	< 65	> 65	53	18508	[10, 12, 13, 15-17, 19-24, 26-30, 32, 34, 46-48, 55, 58, 59, 63, 68, 69, 71-83, 85-90, 92-96]
Mentzer [97]	$MCV / RBC$	< 13	> 13	78	26704	[10-30, 32-36, 39-41, 43-48, 50, 55, 57-61, 63-68, 70-90, 92, 98-103]
RBC [36]	RBC	< 5	> 5	44	11896	[10, 12-14, 16, 18-21, 26, 28, 29, 36, 38-41, 56, 58-60, 65, 67-69, 71-78, 80-83, 86, 88, 99, 103-105]
Bessman (RDW) [106]	RDW	< 14	> 14	52	15208	[10, 12-14, 16, 17, 19-21, 26, 28, 39, 41, 44-48, 50-52, 56-59, 61, 63, 68, 69, 71-77, 80, 82, 83, 86, 88, 94, 98, 100, 104, 105, 107-111]
Jayabose (RDWI) [99]	$(MCV \times RDW) / RBC$	< 220	> 220	34	13916	[10, 12, 13, 15-17, 19-21, 26-30, 68, 71-73, 75-83, 85-88, 92, 96, 112]
Ricerca [98]	$RDW / RBC$	< 4.4	> 4.4	30	13639	[10, 12, 13, 15-17, 19-23, 26, 29, 32, 59, 71, 72, 74, 75, 78-80, 82, 83, 85-89]
Shine & Lal (S&L) [113]	$MCV^2 \times MCH \times 0.01$	<15 30	>15 30	56	21685	[10-17, 19-23, 25-30, 32, 33, 39, 40, 43, 45-48, 50, 55, 60, 61, 65-68, 70-72, 74-76, 78-80, 82, 84-87, 89, 92, 93]
Sirdah [17]	$MCV - RBC - 3 \times Hb$	< 27	> 27	18	9958	[10-15, 17, 19-23, 25-30]
Srivastava [114]	$MCH / RBC$	< 3.8	> 3.8	53	20257	[10-30, 32, 33, 36, 39, 45, 46, 55, 59, 60, 63-65, 67, 68, 71, 72, 74-76, 78-80, 82, 83, 85-89, 92, 98]
M/H Ratio [32]	$\text{Microcytic RBC \%} / \text{hypochromic RBC \%}$	< 3.7	> 3.7	15	3091	[22, 32-34, 59, 63, 89, 90, 103, 115-119]

### Material and Methods

This systematic review and meta-analysis was designed according to the PRISMA guideline [120]. The search strategy in this study was based on the search strategy used by Hoffmann et al. [112].

#### Literature search

To find relevant published papers, we searched six international indexing databases, including PubMed/Medline, ISI web of science, Cochrane

central, ProQuest, Embase, and Scopus up to Dec 2018 systematically. We included the following search terms in the analyses: "microcytic", "iron deficiency", "anemia" or "anaemia" or "iron-deficiency" and/or "thalassemia" AND "distinguish or differentiate or discriminant", and all possible combinations. We did not restrict ourselves regarding the language of the reports and occasionally used the help of a translator.

### **Inclusion criteria**

Two authors (FR and MJ) independently did the initial screening (title and abstract) for inclusion. We identified the original publications of all discriminant indices and searched for the publications citing them. Finally, we perused the literature reference lists in the publications found above for references not yet covered. Other publication types such as the letter, opinion, systematic review or meta-analysis etc. were excluded.

### **Quality assessment**

The quality of published papers was assessed independently by two authors (FR and ASM) using two 12-item and 3-item tools called the Quality Assessment of Diagnostic Accuracy Studies (QUADAS) [121], and the Standards for Reporting Studies of Diagnostic Accuracy (STARD), respectively [122].

### **Statistical analysis**

We extracted TP (True Positive), FP (False Positive), FN (False Negative) and TN (True Negative) values from diagnostic 2-by-2 tables of published studies for calculating and pooling accuracy measures such as summary sensitivity (True Positive Rate), specificity (True Negative Rate), positive likelihood ratio (PLR), negative likelihood ratio (NLR), diagnostic odds ratio (DOR) and area under curve (AUC). The sensitivity and specificity are defined as the probability of positive/negative test results in the subjects with/without the disease, respectively. In this study, the sensitivity and specificity present probability of thalassemia trait/iron deficiency anemia test results in the subjects with/without thalassemia trait, respectively.

### **PLR**

PLR indicates that how many times more likely positive (thalassemia trait) test results in the subjects with the disease (subjects with thalassemia trait) versus the subjects without the disease (subjects with iron deficiency anemia). Also, NPR describes that how many times more likely negative (iron deficiency anemia) test results in the subjects with the disease (subjects with thalassemia trait) versus the subjects without the disease (subjects with iron deficiency anemia).  $PLR > 10$  and  $NLR < 0.1$  indicate a good differential diagnosis [123]. DOR (PLR/NLR) is equal to the ratio of the odds of

positivity (thalassemia trait) in the subjects with disease (thalassemia trait) relative to the odds positivity (thalassemia trait) in the subjects without the disease (iron deficiency anemia) and discriminant index with higher DOR value show better diagnostic differential performance [124]. SROC (summary receiver operating characteristic) curve provides AUC as a global summary of discriminant index accuracy in discriminating between thalassemia trait and iron deficiency anemia and is drawn for each discriminant index. AUC represents an overall performance measure of diagnostic classification and good diagnostic tests have  $AUC > 0.70$  [123].

The heterogeneity across studies was assessed using inconsistency index ( $I^2$  statistics) and Cochran Q test (Chi-square). An  $I^2$  statistic  $> 50\%$  or  $P < 0.1$  for Q test show substantial or significant heterogeneity among studies [125, 126]. If the heterogeneity among the studies was not substantial or significant, pooled summary accuracy would measure with their 95% confidence interval computed based on the fixed-effect model otherwise, the random-effect model is used [127, 128]. The bivariate generalized linear mixed model was also fitted to compare sensitivity and false positive rate of discriminant indices. By using a random effects model, this bivariate regression model incorporates any correlation between sensitivity and false positive rate of different discriminant indices [129]. Publication bias was assessed using Deeks' funnel plot and Deeks' test and a  $P < 0.05$  indicates the presence of publication bias [130]. Data analysis was performed using two software: Meta-Disc version 1.4 (XI Cochrane Colloquium, Barcelona, Spain) [131] and R 3.0.3.

### **Results**

A systematic literature search returned 102 articles reported the usage of 12 different indices that were investigated five or more times (Figure 1). The SROC curves with confidence region and AUCs of all discriminant indices have been measured (Supplementary).

Deeks' tests of all discriminant indices indicated that there is no potential publication bias (Table 2) and Deeks' funnel plots of all discriminant indices did not show any evidence of asymmetry (Deeks' funnel plots are not shown). SROC curves with confidence region are shown in Figure 2 and AUCs of all

## 19 Diagnosis of Microcytic Anemia, Thalassemia or Iron Deficiency Anemia....

discriminant indices indicate overall good differential performance (Table 2). Consequently, the M/H ratio index has higher AUC in comparison with other indices. Bivariate regression model using Bessman (red cell distribution width = RDW) index

as the reference category indicated that sensitivity of all indices is significantly higher than the sensitivity of Bessman (RDW) index ( $P < 0.05$ ).




Fig. 1. The study selection flow diagram

But the false positive rate of Bessman (RDW) index is significantly lower than Ricerca and Shine & Lal indices ( $P < 0.001$ ) and also false positive rate of Bessman (RDW) index is lower than Srivastava index, but there was no statistically significant difference between the false positive rate of Bessman (RDW) index and Srivastava index ( $P > 0.05$ ) (Table 3). The false positive rate of England &

Fraser and Sirdah indices were significantly lower than Bessman (RDW) index ( $P < 0.05$ ) and false positive rate of Ehsani, Green & King, Mentzer, RBC and Jayabose (RDWI) indices were lower than Bessman (RDW) index, but there was no statistically significant difference between the false positive rate of these indices and Bessman (RDW) index ( $P > 0.05$ ) (Table 3).

*Table 2. Diagnostic performance of discriminant indices with 95% confidence interval*

Discriminant Index	Sensitivity (95% CI)	Specificity (95% CI)	PLR (95% CI)	NLR (95% CI)	DOR (95% CI)	AUC	Deeks' Test
Ehsani [9]	0.82 (0.81,0.83)	0.81 (0.80,0.83)	4.57 (3.66,5.71)	0.19 (0.14,0.25)	27.93 (19.23,40.57)	0.905	0.87
England & Fraser (E&F) [31]	0.77 (0.76,0.78)	0.85 (0.84,0.86)	7.71 (6.14,9.7)	0.27 (0.20,0.37)	33.67 (24.83,45.65)	0.9135	0.26
Green & King (G&K) [91]	0.81 (0.80,0.82)	0.85 (0.84,0.86)	6.49 (4.91,8.59)	0.2 (0.14,0.28)	35.35 (23.57,53.02)	0.919	0.82
Mentzer [97]	0.81 (0.81,0.82)	0.83 (0.82,0.84)	4.76 (4.12,5.50)	0.21 (0.18,0.26)	25.68 (19.95,33.08)	0.906	0.62
RBC [36]	0.83 (0.81,0.84)	0.83 (0.82,0.84)	6.14 (4.43,8.49)	0.2 (0.15,0.27)	35.46 (23.49,53.53)	0.921	0.58
Bessman (RDW) [106]	0.63 (0.62,0.64)	0.63 (0.62,0.65)	2.16 (1.76,2.65)	0.45 (0.33,0.62)	6.1 (3.63,10.27)	0.808	0.56
Jayabose (RDWI) [99]	0.84 (0.83,0.85)	0.84 (0.83,0.85)	5.13 (3.7,7.11)	0.2 (0.15,0.25)	28.91 (20.07,41.64)	0.912	0.39
Ricerca [98]	0.89 (0.89,0.90)	0.57 (0.56,0.59)	2.28 (1.79,2.89)	0.17 (0.12,0.22)	19.89 (14.07,28.12)	0.899	0.73
Shine & Lal (S&L) [113]	0.90 (0.90,0.91)	0.52 (0.51,0.53)	1.90 (1.55,2.32)	0.17 (0.13,0.23)	16.52 (10.40,26.24)	0.899	0.61
Sirdah [17]	0.79 (0.77,0.80)	0.86 (0.85,0.87)	7.71 (5.73,10.38)	0.25 (0.19,0.34)	39.30 (25.90, 59.64)	0.933	0.70
Srivastava [114]	0.77 (0.76,0.78)	0.79 (0.78,0.80)	3.85 (3.27,4.52)	0.31 (0.26,0.36)	14.25 (10.92,18.60)	0.861	0.57
M/H Ratio [32]	0.92 (0.87,0.98)	0.86 (0.81,0.91)	6.8 (4.8,9.8)	0.07 (0.03,0.2)	100.8 (39.6,256.3)	0.956	


CI: Confidence Interval, PLR: Positive Likelihood Ratio; NLR: Negative Likelihood Ratio; DOR: Diagnostic Odds Ratio,; AUC: Area Under Curve.

*Table 3. The result of analysis of data by bivariate generalized linear mixed model*

Discriminant Index	Sensitivity			False Positive Rate		
	Estimate	95% CI	P-value	Estimate	95% CI	P-value
Intercept	0.201	(-0.217,0.619)	0.346	-1.526	(-1.967,-1.085)	<0.001
Ehsani	1.566	(0.915,2.217)	<0.001	-0.013	(-0.683,0.658)	0.970
England & Fraser (E&F)	0.810	(0.289,1.331)	0.002	-0.862	(-1.418,-0.306)	0.002
Green & King (G&K)	1.310	(0.767,1.854)	<0.001	-0.387	(-0.958,0.184 )	0.184
Mentzer	1.333	(0.816,1.849)	<0.001	-0.134	(-0.676,0.408)	0.628
RBC	1.377	(0.809,1.945)	<0.001	-0.318	(-0.918,0.281)	0.298
Jayabose (RDWI)	1.496	(0.910,2.082)	<0.001	-0.047	(-0.654,0.559)	0.878
Ricerca	2.374	(1.750,2.998)	<0.001	1.370	(0.740,2.001)	<0.001
Shine & Lal (S&L)	2.669	(2.097,3.242)	<0.001	1.886	(1.314,2.458)	<0.001
Sirdah	1.069	(0.388,1.750)	0.002	-0.782	(-1.495,-0.069 )	0.032
Srivastava	0.916	(0.379,1.453)	0.001	0.002	(-0.565,0.568)	0.995
M/H Ratio	1.8	(1.1,2.5)	<0.001	-0.6	(-1.3, 0.1)	0.1

Bessman (RDW) index is used as reference group. CI: Confidence Interval.

## 21 Diagnosis of Microcytic Anemia, Thalassemia or Iron Deficiency Anemia....



*Fig. 2. SROC (summary receiver operating characteristic) curves with confidence region of discriminant indices.*

## Discussion

Microcytic anemia is one of the most common forms of anemia that physicians deal with during *daily practice*, and are often faced with the challenge of its managing. The differentiation of the two most common forms of microcytic anemia which include IDA and  $\beta$ -thalassemia minor is very important because, in addition to high prevalence, it is sometimes difficult to differentiate them; therefore patients suffer from loss of time and money. Today, screening of thalassemia minor has a special place in countries dealing with a huge number of families at risk of a severe form of the disease. Identifying minor thalassemia is also important because the treatment of these patients with iron leads to increase in mean corpuscular volume (MCV) levels. The main method of anemia diagnosis is a reduction in serum iron level, saturated transferrin, and ferritin levels with an increase in iron binding capacity. Although, the method of thalassemia minor diagnosis is measuring HbA2 levels by means of hemoglobin electrophoresis test [132], in some cases of thalassemia minor, HbA2 levels do not rise and on the other hand, measurement of HbA2 levels or iron storages in the body are considered as time-consuming and costly methods. Moreover, there is a controversy regarding whether HbA2 levels provide a valid and *reliable measure* of diagnosis of  $\beta$ -thalassemia trait in people with iron deficiency or not [133].

Nowadays, other low-cost and rapid calculation methods based on the measurement of the red blood cell indices are very common and have been used as first-line diagnostic methods so far. The sensitivity and specificity of these methods have been assessed in various studies and their validity has been proven [134-139]. Since 1973, there has always been an attempt to introduce newer and more efficient discriminating indices between IDA and thalassemia minor. The calculations of these indices are sometimes so easy and sometimes challenging. The dissimilarity of age and sex groups and the difference in serum hemoglobin and ferritin thresholds in detecting IDA or thalassemia could lead to categorizing patients with various severities of iron deficiency as an IDA group [140].

IDA and thalassemia trait RBC indices resemblance in CBC test makes it hard to only lean on CBC test in discriminating them. On the other hand, performing hemoglobin electrophoresis and more importantly, the interpretation of its results are not applicable in every thalassemia endemic region. These are some of the reasons why the usage of mathematical indices is beneficial and useful in discriminating IDA from thalassemia trait even with their lack of full sensitivity and specificity [112]. Moreover, it is necessary to have in mind that previous studies evaluating these indices may have not completely considered some of the effective parameters which cause a major limitation in the study of mathematical indices. For instance, how to introduce cutoff values, study design, patient inclusion criteria and thalassemia genotype of patients are all among affective parameters which are required to be considered in order to reach a definite outcome in the analysis. Among introduced indices, it seems that later ones are more reliable in discriminating IDA from thalassemia trait because of their higher sensitivity and specificity.

## Conclusion and clinical application

In this meta-analysis, we showed that the M/H ratio index is more sensitive and specific compared to our other studied indices. Therefore, it has more potential to discriminate IDA from thalassemia trait. However, we still cannot use this index alone to achieve a final diagnosis. The capability of this index to discriminate IDA from thalassemia trait must be used alongside with standard guidelines for identification of these entities to ensure the final differentiation. Ultimately, the detection of IDA from thalassemia trait using these indices is a valuable diagnostic approach. Although these tests have relatively high sensitivity and specificity, the result should be interpreted based on the combination of clinical and laboratory ratifications. Further studies regarding the measurement of the biomarkers for screening and fast detection are suggested.

## Conflict of Interest

Authors declared no conflict of interest.

## Acknowledgments

We are grateful to all colleagues who help us in the study.

## 23 Diagnosis of Microcytic Anemia, Thalassemia or Iron Deficiency Anemia....

### References

1. DeLoughery TG. Microcytic anemia. *The New England journal of medicine*. 2014;371(14):1324-31.
2. Stuckey WJ. Common anemias: a practical guide to diagnosis and management. *Geriatrics*. 1983;38(8):42-8.
3. Camaschella C. New insights into iron deficiency and iron deficiency anemia. *Blood reviews*. 2017;31(4):225-33.
4. Li C-K. New trend in the epidemiology of thalassaemia. *Best Practice & Research Clinical Obstetrics & Gynaecology*. 2017;39:16-26.
5. DeLoughery TG. Microcytic anemia. *New England Journal of Medicine*. 2014;371(14):1324-31.
6. Barrett AN, Saminathan R, Choolani M. Thalassaemia screening and confirmation of carriers in parents. *Best Practice & Research Clinical Obstetrics & Gynaecology*. 2017;39:27-40.
7. Ntaios G, Chatzinikolaou A. Discrimination indices as screening tests for  $\beta$ -thalassemic trait. *Annals of Hematology*. 2008;87(4):329-30.
8. Mentzer WC, Jr. Differentiation of iron deficiency from thalassaemia trait. *Lancet (London, England)*. 1973;1(7808):882.
9. Ehsani M, Shahgholi E, Rahiminejad M, Seighali F, Rashidi A. A new index for discrimination between iron deficiency anemia and beta-thalassemia minor: results in 284 patients. *Pakistan journal of biological sciences: PJBS*. 2009;12(5):473-5.
10. Chandra H, Shrivastava V, Chandra S, Rawat A, Nautiyal R. Evaluation of Platelet and Red Blood Cell Parameters with Proposal of Modified Score as Discriminating Guide for Iron Deficiency Anemia and  $\beta$ -Thalassemia Minor. *Journal of clinical and diagnostic research: JCDR*. 2016;10(5):EC31.
11. Bordbar E, Taghipour M, Zucconi BE. Reliability of different RBC indices and formulas in discriminating between  $\beta$ -thalassemia minor and other microcytic hypochromic cases. *Mediterranean journal of hematology and infectious diseases*. 2015;7(1).
12. Urrechaga E, Hoffmann JJ. Critical appraisal of discriminant formulas for distinguishing thalassemia from iron deficiency in patients with microcytic anemia. *Clinical Chemistry and Laboratory Medicine (CCLM)*. 2017.
13. Pornprasert S, Thongsat C, Panyachadporn U. Evaluation of Applying a Combination of Red Cell Indexes and Formulas to Differentiate  $\beta$ -Thalassemia Trait from Iron Deficiency Anemia in the Thai Population. *Hemoglobin*. 2017:1-4.
14. Susanti AI, Sahiratmadja E, Winarno G, Sugianli AK, Susanto H, Panigoro R. Low Hemoglobin among Pregnant Women in Midwives Practice of Primary Health Care, Jatinangor, Indonesia: Iron Deficiency Anemia or  $\beta$ -Thalassemia Trait? *Anemia*. 2017;2017.
15. Huang T-C, Wu Y-Y, Chen Y-G, Lai S-W, Wu S-C, Ye R-H, et al. Discrimination index of microcytic anemia in young soldiers: a single institutional analysis. *PloS one*. 2015;10(2):e0114061.
16. Zaghoul A, Al-bukhari T, Bajuaifer N, Shalaby M, Al-Pakistani H, Halawani SH, et al. Introduction of new formulas and evaluation of the previous red blood cell indices and formulas in the differentiation between beta thalassemia trait and iron deficiency anemia in the Makkah region. *Hematology*. 2016;21(6):351-8.
17. Sirdah M, Tarazi I, Al Najjar E, Al Haddad R. Evaluation of the diagnostic reliability of different RBC indices and formulas in the differentiation of the  $\beta$ -thalassaemia minor from iron deficiency in Palestinian population. *International Journal of Laboratory Hematology*. 2008;30(4):324-30.
18. Ehsani M, Shahgholi E, Rahiminejad M, Seighali F, Rashidi A. A new index for discrimination between iron deficiency anemia and beta-thalassemia minor: results in 284 patients. *Pakistan Journal of Biological Sciences*. 2009;12(5):473-5.
19. Keikhaei B. A new valid formula in differentiating iron deficiency anemia from  $\beta$ -thalassemia trait. *Pakist J Med Sci*. 2010;26:368-73.
20. Janel A, Roszyk L, Rapatel C, Mareynat G, Berger MG, Serre-Sapin AF. Proposal of a score combining red blood cell indices for early differentiation of beta-thalassemia minor from iron deficiency anemia. *Hematology*. 2011;16(2):123-7.
21. Shen C, Jiang Y-m, Shi H, Liu J-h, Zhou W-j, Dai Q-k, et al. Evaluation of indices in differentiation between iron deficiency anemia and  $\beta$ -thalassemia

- trait for Chinese children. *Journal of pediatric hematology/oncology*. 2010;32(6):e218-e22.
22. Urrechaga E, Borque L, Escanero J. The role of automated measurement of red cell subpopulations on the Sysmex XE 5000 analyzer in the differential diagnosis of microcytic anemia. *International Journal of Laboratory Hematology*. 2011;33(1):30-6.
  23. Urrechaga E, Borque L, Escanero JF. The role of automated measurement of RBC subpopulations in differential diagnosis of microcytic anemia and  $\beta$ -thalassemia screening. *American journal of clinical pathology*. 2011;135(3):374-9.
  24. Nishad AAN, Pathmeswaran A, Wickremasinghe A, Premawardhana A. The Thal-index with the BTT prediction. *exe to discriminate  $\beta$ -thalassaemia traits from other microcytic anaemias*. 2012.
  25. Sudmann Å, Piehler A, Urdal P. Reticulocyte hemoglobin equivalent to detect thalassemia and thalassemic hemoglobin variants. *International Journal of Laboratory Hematology*. 2012;34(6):605-13.
  26. Sahli CA, Bibi A, Ouali F, Fredj SH, Dakhlaoui B, Othmani R, et al. Red cell indices: differentiation between  $\beta$ -thalassemia trait and iron deficiency anemia and application to sickle-cell disease and sickle-cell thalassemia. *Clinical chemistry and laboratory medicine*. 2013;51(11):2115-24.
  27. Miri-Moghaddam E, Sargolzaie N. Cut off determination of discrimination indices in differential diagnosis between iron deficiency anemia and  $\beta$ -thalassemia minor. *International journal of hematology-oncology and stem cell research*. 2014;8(2):27.
  28. Pornprasert S, Panya A, Punyamung M, Yanola J, Kongpan C. Red cell indices and formulas used in differentiation of  $\beta$ -thalassemia trait from iron deficiency in Thai school children. *Hemoglobin*. 2014;38(4):258-61.
  29. Vehapoglu A, Ozgurhan G, Demir AD, Uzuner S, Nursoy MA, Turkmen S, et al. Hematological indices for differential diagnosis of beta thalassemia trait and iron deficiency anemia. *Anemia*. 2014;2014.
  30. Matos JF, Dusse LMSA, Stubbart RVB, Ferreira MR, Coura-Vital W, Fernandes APSM, et al. Comparison of discriminative indices for iron deficiency anemia and  $\beta$  thalassemia trait in a Brazilian population. *Hematology*. 2013;18(3):169-74.
  31. En gland J, Fraser P. Differentiation of iron deficiency from thalassaemia trait by routine blood-count. *The Lancet*. 1973;301(7801):449-52.
  32. Ur rechaga E. Discriminant value of% microcytic/% hypochromic ratio in the differential diagnosis of microcytic anemia. *Clinical chemistry and laboratory medicine*. 2008;46(12):1752-8.
  33. Vi cinanza P, Catalano L, Franco F, Vaccaro E, Cancellario S, Vicinanza M, et al. Two New HDW-Based Indexes to Identify beta-Thalassemia Trait. *Laboratory Hematology*. 2002;8:193-9.
  34. Sc hoorl M, Schoorl M, Linszen J, Villanueva MM, NoGuera JAV, Martinez PH, et al. Efficacy of advanced discriminating algorithms for screening on iron-deficiency anemia and  $\beta$ -thalassemia trait: a multicenter evaluation. *American journal of clinical pathology*. 2012;138(2):300-4.
  35. Gi mferre E, Marigo G, Rutllant M, VIÑAS J. Differentiation of iron deficiency from thalassaemia trait. *The Lancet*. 1975;305(7898):114.
  36. Kl ee GG, Fairbanks VF, Pierre RV, O'sullivan MB. Routine erythrocyte measurements in diagnosis of iron-deficiency anemia and thalassemia minor. *American journal of clinical pathology*. 1976;66(5):870-7.
  37. He gde U, White J, Hart G, Marsh G. Diagnosis of alpha-thalassemia trait from Coulter Counter'S'indices. *Journal of clinical pathology*. 1977;30(9):884-9.
  38. Wa lford D, McPherson K, Deacon R. Discrimination between iron deficiency and heterozygous thalassaemia. *The Lancet*. 1979;313(8111):323.
  39. Jo hnson CS, Tegos C, Beutler E. Thalassemia Minor: Routine Erythrocyte Measurements and Differentiation from Iron Deficienc. *American journal of clinical pathology*. 1983;80(1):31-6.
  40. Ch alevelakis G, Tsiroyannis K, Hatzioannou J, Arapakis G. Screening for thalassaemia and/or iron deficiency: evaluation of some discrimination functions. *Scandinavian journal of clinical and laboratory investigation*. 1984;44(1):1-6.
  41. Gh ionni H, Miotti T, Camandona V. Routine erythrocyte measurements and differentiation of

## 25 Diagnosis of Microcytic Anemia, Thalassemia or Iron Deficiency Anemia....

- thalassemia minor from iron deficiency. *Minerva Med.* 1985;76:1143-8.
42. Juncà PJ, Clotet SB, Millà SF, Ribas MM. England's index in microcytosis. *Medicina clinica.* 1985;85(18):773-.
  43. Marti HR, Fischer S, Killer D, Bürgi W. Can automated haematology analysers discriminate thalassaemia from iron deficiency? *Acta haematologica.* 1987;78(2-3):180-3.
  44. Novak RW. Red blood cell distribution width in pediatric microcytic anemias. *Pediatrics.* 1987;80(2):251-4.
  45. Helleman P, Bartels P, van Waveren Hogervorst G. Screening for thalassaemia using the width of the Technicon H6000/H601® erythrocyte size histograms. *Scandinavian journal of clinical and laboratory investigation.* 1988;48(7):697-704.
  46. Bentley S, Ayscue L, Watson J, Ross D. The clinical utility of discriminant functions for the differential diagnosis of microcytic anemias. *Blood cells.* 1989;15(3):575-82; discussion 83-4.
  47. Houwen B. The use of inference strategies in the differential diagnosis of microcytic anemia. *Blood cells.* 1989;15(3):509-27; discussion 27-32.
  48. Makris P. Utilization of a new index to distinguish heterozygous thalassemic syndromes: comparison of its specificity to five other discriminants. *Blood cells.* 1989;15(3):497-506; discussion 7.
  49. Paterakis G, Terzoglou G, Vasilioy E. The performance characteristics of an expert system for the " on-line" assessment of thalassemia trait and iron deficiency--Micro Hema Screen. *Blood cells.* 1989;15(3):541-60; discussion 60-1.
  50. Qurtom H, Al-Saleh Q, Lubani M, Hassanein A, Kaddoorah N, Qurtom M, et al. The value of red cell distribution width in the diagnosis of anaemia in children. *European journal of pediatrics.* 1989;148(8):745-8.
  51. Laso F, Mateos F, Ramos R, Herrero F, Pérez-Arellano J, González-Buitrago J. Amplitude of the distribution of erythrocyte size in the differential diagnosis of microcytic anemia. *Medicina clinica.* 1990;94(1):1-4.
  52. Junca J, Flores A, Roy C, Alberti R, Milla F. Red cell distribution width, free erythrocyte protoporphyrin, and England-Fraser index in the differential diagnosis of microcytosis due to iron deficiency or beta-thalassemia trait. A study of 200 cases of microcytic anemia. *Hematologic pathology.* 1991;5(1):33-6.
  53. Ortega CA, Pineda AM, Pardo AJ, Sánchez RJ. The prognostic value of the hemogram in the differential diagnosis of microcytic anemia. *Medicina clinica.* 1991;96(10):397.
  54. Robertson E, Pollock A, Yau K, Chan L. Use of technicon H\* 1 technology in routine thalassaemia screening. *Medical laboratory sciences.* 1992;49(4):259-64.
  55. Jiménez CV. Iron-deficiency anemia and thalassemia trait differentiated by simple hematological tests and serum iron concentrations. *Clinical chemistry.* 1993;39(11):2271-5.
  56. Baqar MS, Khurshid M, Molla A. Does red blood cell distribution width (RDW) improve evaluation of microcytic anaemias? *Journal of Pakistan Medical Association.* 1993;43(8):149.
  57. Das GA, Hegde C, Mistri R. Red cell distribution width as a measure of severity of iron deficiency in iron deficiency anaemia. *The Indian journal of medical research.* 1994;100:177-83.
  58. Burk M, Arenz J, Giagounidis A, Schneider W. Erythrocyte indices as screening tests for the differentiation of microcytic anemias. *European journal of medical research.* 1995;1(1):33-7.
  59. Erler BS, Vitagliano P, Lee S. Superiority of neural networks over discriminant functions for thalassemia minor screening of red blood cell microcytosis. *Archives of pathology & laboratory medicine.* 1995;119(4):350-4.
  60. Yeo G, Tan K, Liu T. The role of discriminant functions in screening for beta-thalassaemia traits during pregnancy. *Singapore medical journal.* 1995;36(6):615-8.
  61. Lafferty JD, Crowther MA, Ali MA, Levine M. The evaluation of various mathematical RBC indices and their efficacy in discriminating between thalassemic and non-thalassemic microcytosis. *American journal of clinical pathology.* 1996;106(2):201-5.
  62. Mach-Pascual S, Darbellay R, Pilotto PA, Beris P. Investigation of microcytosis: a comprehensive approach. *European journal of haematology.* 1996;57(1):54-61.
  63. Liu T-C, Seong P-S, Lin T-K. The erythrocyte cell hemoglobin distribution width segregates thalassemia traits from other nonthalassemic

- conditions with microcytosis. *American journal of clinical pathology*. 1997;107(5):601-7.
64. Manglani M, Lokeshwar M, Vani V, Bhatia N, Mhaskar V. 'NESTROFT'--an effective screening test for beta thalassemia trait. *Indian pediatrics*. 1997;34(8):702-7.
  65. Madan N, Sikka M, Sharma S, Rusia U, Kela K. Red cell indices and discriminant functions in the detection of beta-thalassaemia trait in a population with high prevalence of iron deficiency anaemia. *Indian journal of pathology & microbiology*. 1999;42(1):55-61.
  66. TELMISSANI OA, KHALIL S, ROBERTS GT. Mean density of hemoglobin per liter of blood: a new hematologic parameter with an inherent discriminant function. *Laboratory Hematology*. 1999;5:149-52.
  67. Kuznetsova Y, Kovrigina Y, Baidun L, Bykova L, Chernov V, Tokarev Y, et al. Use of erythrocytic indexes and iron metabolism parameters in the differential diagnosis of microcytic anemias. *GEMATOLOGIYA I TRANSFUZIOLOGIYA*. 2000;45(6):46-8.
  68. Demir A, Yarali N, Fisgin T, Duru F, Kara A. Most reliable indices in differentiation between thalassemia trait and iron deficiency anemia. *Pediatrics International*. 2002;44(6):612-6.
  69. Melo MR, Purini MC, Caçado RD, Koro F, Chiattoni CS. The use of erythrocyte (RBC) indices in the differential diagnosis of microcytic anemias: is it an approach to be adopted? *Revista da Associacao Medica Brasileira (1992)*. 2002;48(3):222-4.
  70. Ghafouri M, MOSTAAN SL, Sharifi S, HOSSEINI GL, ATAR CZ. Comparison of cell counter indices in differentiation of beta thalassemia minor from iron deficiency anemia. 2006.
  71. AlFadhli SM, Al-Awadhi AM, AlKhalidi Da. Validity assessment of nine discriminant functions used for the differentiation between iron deficiency anemia and thalassemia minor. *Journal of tropical pediatrics*. 2006;53(2):93-7.
  72. Beyan C, Kaptan K, Ifran A. Predictive value of discrimination indices in differential diagnosis of iron deficiency anemia and beta-thalassemia trait. *European journal of haematology*. 2007;78(6):524-6.
  73. Ntaios G, Chatzinikolaou A, Saouli Z, Girtovitis F, Tsapanidou M, Kaiafa G, et al. Discrimination indices as screening tests for  $\beta$ -thalassemic trait. *Annals of hematology*. 2007;86(7):487-91.
  74. Rathod DA, Kaur A, Patel V, Patel K, Kabrawala R, Patel V, et al. Usefulness of cell counter-based parameters and formulas in detection of  $\beta$ -thalassemia trait in areas of high prevalence. *American journal of clinical pathology*. 2007;128(4):585-9.
  75. Okan V, Cigiloglu A, Cifci S, Yilmaz M, Pehlivan M. Red cell indices and functions differentiating patients with the  $\beta$ -thalassaemia trait from those with iron deficiency anaemia. *Journal of International Medical Research*. 2009;37(1):25-30.
  76. Rahim F, Keikhaei B. Better differential diagnosis of iron deficiency anemia from beta-thalassemia trait. *Turk J Hematol*. 2009;26(3):138-45.
  77. Ferrara M, Capozzi L, Russo R, Bertocco F, Ferrara D. Reliability of red blood cell indices and formulas to discriminate between  $\beta$  thalassemia trait and iron deficiency in children. *Hematology*. 2010;15(2):112-5.
  78. Narchi H, Basak R. Comparison of erythrocyte indices to differentiate between iron deficiency and alpha-thalassaemias in children with microcytosis an. 2010.
  79. Niazi M, Tahir M, e Raziq F, Hameed A. Usefulness of Redcell Indices in Differentiating Microcytic Hypochromic Anemias. *Gomal Journal of Medical Sciences*. 2010;8(2).
  80. Tiwari A, Chandola I, Ahuja A. Approach to blood donors with microcytosis. *Transfusion Medicine*. 2010;20(2):88-94.
  81. Nesa A, Munir S, Sultana T, Rahman M, Ahmed A. Role of discrimination indices in differentiation of beta thalassaemia trait and iron deficiency anaemia. *Mymensingh medical journal: MMJ*. 2011;20(1):110-4.
  82. Trivedi D, Shah H. Discriminant functions in distinguishing beta-thalassemia trait and iron deficiency anemia: the value of the RDW-SD. *Internet J Hematol*. 2011;7:1-13.
  83. Williams C. The use of the automated parameter RDW-SD as a discriminator of both alpha and beta thalassaemia trait from iron deficiency. *Australian Journal of Medical Science*. 2011;32(3):88.

## 27 Diagnosis of Microcytic Anemia, Thalassemia or Iron Deficiency Anemia....

84. Batebi A, Pourreza A, Esmailian R. Discrimination of beta-thalassemia minor and iron deficiency anemia by screening test for red blood cell indices. *Turkish Journal of Medical Sciences*. 2012;42(2):275-80.
85. Gibson F, Mason K, Serjeant B, Kulozik A, Happich M, Tolle G, et al. Screening for the beta-thalassaemia trait: hazards among populations of West African Ancestry. *Journal of community genetics*. 2012;3(1):13-8.
86. Nalbantoğlu B, Güzel S, Büyükyalçın V, Donma MM, Güzel EÇ, Nalbantoğlu A, et al. Indices used in differentiation of thalassemia trait from iron deficiency anemia in pediatric population: are they reliable? *Pediatric hematology and oncology*. 2012;29(5):472-8.
87. Dharmani P, Sehgal K, Dadu T, Mankeshwar R, Shaikh A, Khodaiji S. Developing a new index and its comparison with other CBC-based indices for screening of beta thalassemia trait in a tertiary care hospital. *International Journal of Laboratory Hematology*. 2013;35:118.
88. Ng E, Leung J, Lau Y, Ma E. Evaluation of the new red cell parameters on Beckman Coulter DxH800 in distinguishing iron deficiency anaemia from thalassaemia trait. *International Journal of Laboratory Hematology*. 2015;37(2):199-207.
89. Urrechaga E, Hoffmann J, Izquierdo S, Escanero J. Differential diagnosis of microcytic anemia: the role of microcytic and hypochromic erythrocytes. *International Journal of Laboratory Hematology*. 2015;37(3):334-40.
90. Yuen S, Brown R, Forsyth C, Zarkos K. A quick differentiation between iron deficiency and thalassemia syndrome. *Australian Journal of Medical Science*. 1996;17:106-9.
91. Green R, King R. A new red cell discriminant incorporating volume dispersion for differentiating iron deficiency anemia from thalassemia minor. *Blood cells*. 1989;15(3):481-95.
92. Ullah Z, Khattak AA, Ali SA, Hussain J, Noor B, Bano R, et al. Evaluation of five discriminating indexes to distinguish Beta-Thalassemia Trait from Iron Deficiency Anaemia. *JPMA The Journal of the Pakistan Medical Association*. 2016;66(12):1627.
93. Bessman J, McClure S, Bates J. Distinction of microcytic disorders: comparison of expert, numerical-discriminant, and microcomputer analysis. *Blood cells*. 1989;15(3):533-40.
94. Lima CSP, Reis ARdC, Grotto HZW, Saad STO, Costa FF. Comparison of red cell distribution width and a red cell discriminant function incorporating volume dispersion for distinguishing iron deficiency from beta thalassemia trait in patients with microcytosis. *Sao Paulo Medical Journal*. 1996;114(5):1265-9.
95. Nadarajan V, Sthaneshwar P, Jayarane S. RBC-Y/MCV as a discriminant function for differentiating carriers of thalassaemia and HbE from iron deficiency. *International Journal of Laboratory Hematology*. 2010;32(2):215-21.
96. Wongprachum K, Sanchaisuriya K, Sanchaisuriya P, Siridamrongvattana S, Manpeun S, Schlep FP. Proxy indicators for identifying iron deficiency among anemic vegetarians in an area prevalent for thalassemia and hemoglobinopathies. *Acta haematologica*. 2012;127(4):250-5.
97. Mentzer W. Differentiation of iron deficiency from thalassaemia trait. *The Lancet*. 1973;301(7808):882.
98. Ricerca B, Storti S, d'Onofrio G, Mancini S, Vittori M, Campisi S, et al. Differentiation of iron deficiency from thalassaemia trait: a new approach. *Haematologica*. 1986;72(5):409-13.
99. Jayabose S, Giamelli J, LevondogluTugal O, Sandoval C, Ozkaynak F, Visintainer P. # 262 Differentiating iron deficiency anemia from thalassemia minor by using an RDW-based index. *Journal of pediatric hematology/oncology*. 1999;21(4):314.
100. Eldibany MM, Totonchi KF, Joseph NJ, Rhone D. Usefulness of certain red blood cell indices in diagnosing and differentiating thalassemia trait from iron-deficiency anemia. *American journal of clinical pathology*. 1999;111(5):676-82.
101. Afroz M, Shamsi TS, Syed S. Predictive Value of MCV/RBC Count Ratio to discriminate between Iron Deficiency Anaemia and Beta Thallasaemia Trait. *JOURNAL-PAKISTAN MEDICAL ASSOCIATION*. 1998;48:18-.
102. Kneifati-Hayek J, Fleischman W, Bernstein LH, Riccioli A, Bellevue R. A model for automated screening of thalassemia in hematology (math study). *Laboratory hematology: official publication of the International Society for Laboratory Hematology*. 2007;13(4):119-23.

103. Urrechaga E. Red blood cell microcytosis and hypochromia in the differential diagnosis of iron deficiency and  $\beta$ -thalassaemia trait. *International Journal of Laboratory Hematology*. 2009;31(5):528-34.
104. Flynn MM, Reppun TS, Bhagavan NV. Limitations of red blood cell distribution width (RDW) in evaluation of microcytosis. *American journal of clinical pathology*. 1986;85(4):445-9.
105. Kotwal J, Saxena R, Choudhry V, Dwivedi S, Bhargava M. Erythrocyte indices for discriminating thalassaemic and non-thalassaemic microcytosis in Indians. *The National medical journal of India*. 1999;12(6):266-7.
106. Bessman JD, Feinstein D. Quantitative anisocytosis as a discriminant between iron deficiency and thalassaemia minor. *Blood*. 1979;53(2):288-93.
107. Miguel A, Linares M, Miguel A, Miguel-Borja JM. Red cell distribution width analysis in differentiation between iron deficiency and thalassaemia minor. *Acta haematologica*. 1988;80(1):59-.
108. Bhamhani K, Aronow R. Lead poisoning and thalassaemia trait or iron deficiency: the value of the red blood cell distribution width. *American Journal of Diseases of Children*. 1990;144(11):1231-3.
109. Zeben D, Bieger R, Wermeskerken RK, Castel A, Hermans J. Evaluation of microcytosis using serum ferritin and red blood cell distribution width. *European journal of haematology*. 1990;44(2):106-9.
110. MAIOLO AT, GIDIULI R, DAMILANO I, MASSARO P, POLLI EE. Relevance of red cell distribution width (RDW) in the differential diagnosis of microcytic anaemias. *International Journal of Laboratory Hematology*. 1991;13(2):141-51.
111. Harrington AM, Ward PC, Kroft SH. Iron deficiency anemia,  $\beta$ -thalassaemia minor, and anemia of chronic disease: a morphologic reappraisal. *American journal of clinical pathology*. 2008;129(3):466-71.
112. Hoffmann JJ, Urrechaga E, Aguirre U. Discriminant indices for distinguishing thalassaemia and iron deficiency in patients with microcytic anemia: a meta-analysis. *Clinical Chemistry and Laboratory Medicine (CCLM)*. 2015;53(12):1883-94.
113. Shine I, Lal S. A strategy to detect  $\beta$ -thalassaemia minor. *The Lancet*. 1977;309(8013):692-4.
114. Srivastava P, Bevington J. Iron deficiency and/or Thalassaemia trait. *The Lancet*. 1973;301(7807):832.
115. d'Onofrio G, Zini G, Ricerca BM, Mancini S, Mango G. Automated measurement of red blood cell microcytosis and hypochromia in iron deficiency and beta-thalassaemia trait. *Archives of pathology & laboratory medicine*. 1992;116(1):84-9.
116. Saleem M, Qureshi T, Anwar M, Ahmed S. Evaluation of M/H ratio for screening of beta thalassaemia trait. 1995.
117. Fossat C, Camoin-Jau L, Marin V, Grob F, David M. Interpretation des anémies chez l'adulte: intérêt des paramètres érythrocytaires. *FEUILLETS DE BIOLOGIE*. 1996:5-12.
118. Kremer A, Kutter D, Groff P. IRON DEFICIENCY OR HAEMOGLOBINOPATHY? DIFFERENTIAL DIAGNOSIS BY RED CELL INDICES PRODUCED BY THE ADVIA BAYER HAEMATOLOGICAL AUTOMAT. *Clinical laboratory*. 1999;45(9-10):547-51.
119. Yermiahu T, Ben-Shalom M, Porath A, Vardi H, Boantz A, Mazor D, et al. Quantitative determinations of microcytic-hypochromic red blood cell population and glycerol permeability in iron-deficiency anemia and  $\beta$  thalassaemia minor. *Annals of hematology*. 1999;78(10):468-71.
120. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *International journal of surgery*. 2010;8(5):336-41.
121. Whiting P, Rutjes AW, Dinnes J, Reitsma J, Bossuyt PM, Kleijnen J. Development and validation of methods for assessing the quality of diagnostic accuracy studies. *Health technology assessment*. 2004;8(25):iii, 1-234.
122. Bossuyt PM, Reitsma JB, Bruns DE, Gatsonis CA, Glasziou PP, Irwig LM, et al. The STARD statement for reporting studies of diagnostic accuracy: explanation and elaboration. *Annals of internal medicine*. 2003;138(1):W1-12.

## 29 Diagnosis of Microcytic Anemia, Thalassemia or Iron Deficiency Anemia....

- 123.Šimundić A-M. Measures of diagnostic accuracy: basic definitions. *EJIFCC*. 2009;19(4):203.
- 124.Glas AS, Lijmer JG, Prins MH, Bossel GJ, Bossuyt PM. The diagnostic odds ratio: a single indicator of test performance. *Journal of clinical epidemiology*. 2003;56(11):1129-35.
- 125.Higgins J, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Statistics in medicine*. 2002;21(11):1539-58.
- 126.Dinnes J, Deeks J, Kirby J, Roderick P. A methodological review of how heterogeneity has been examined in systematic reviews of diagnostic test accuracy. 2005.
- 127.Lee J, Kim KW, Choi SH, Huh J, Park SH. Systematic review and meta-analysis of studies evaluating diagnostic test accuracy: a practical review for clinical researchers-part II. statistical methods of meta-analysis. *Korean journal of radiology*. 2015;16(6):1188-96.
- 128.DerSimonian R, Laird N. Meta-analysis in clinical trials. *Controlled clinical trials*. 1986;7(3):177-88.
- 129.Reitsma JB, Glas AS, Rutjes AW, Scholten RJ, Bossuyt PM, Zwinderman AH. Bivariate analysis of sensitivity and specificity produces informative summary measures in diagnostic reviews. *Journal of clinical epidemiology*. 2005;58(10):982-90.
- 130.Deeks JJ, Macaskill P, Irwig L. The performance of tests of publication bias and other sample size effects in systematic reviews of diagnostic test accuracy was assessed. *Journal of clinical epidemiology*. 2005;58(9):882-93.
- 131.Zamora J, Abaira V, Muriel A, Khan K, Coomarasamy A. Meta-DiSc: a software for meta-analysis of test accuracy data. *BMC medical research methodology*. 2006;6(1):31.
- 132.Joly P, Pondarre C, Badens C. [Beta-thalassemias: molecular, epidemiological, diagnostic and clinical aspects]. *Annales de biologie clinique*. 2014;72(6):639-68.
- 133.Verhovsek M, So CC, O'Shea T, Gibney GT, Ma ES, Steinberg MH, et al. Is HbA2 level a reliable diagnostic measurement for beta-thalassemia trait in people with iron deficiency? *American journal of hematology*. 2012;87(1):114-6.
- 134.Ferrara M, Capozzi L, Russo R, Bertocco F, Ferrara D. Reliability of red blood cell indices and formulas to discriminate between beta thalassemia trait and iron deficiency in children. *Hematology*. 2010;15(2):112-5.
- 135.Nesa A, Munir SF, Sultana T, Rahman MQ, Ahmed AN. Role of discrimination indices in differentiation of beta thalassaemia trait and iron deficiency anaemia. *Mymensingh medical journal : MMJ*. 2011;20(1):110-4.
- 136.Pornprasert S, Thongsat C, Panyachadporn U. Evaluation of Applying a Combination of Red Cell Indexes and Formulas to Differentiate beta-Thalassemia Trait from Iron Deficiency Anemia in the Thai Population. *Hemoglobin*. 2017;41(2):116-9.
- 137.Rahim F, Keikhaei B. Better differential diagnosis of iron deficiency anemia from beta-thalassemia trait. *Turkish journal of haematology : official journal of Turkish Society of Haematology*. 2009;26(3):138-45.
- 138.Sahli CA, Bibi A, Ouali F, Fredj SH, Dakhlaoui B, Othmani R, et al. Red cell indices: differentiation between beta-thalassemia trait and iron deficiency anemia and application to sickle-cell disease and sickle-cell thalassemia. *Clin Chem Lab Med*. 2013;51(11):2115-24.
- 139.Zaghloul A, Al-Bukhari TA, Bajuaifer N, Shalaby M, Al-Pakistani HA, Halawani SH, et al. Introduction of new formulas and evaluation of the previous red blood cell indices and formulas in the differentiation between beta thalassemia trait and iron deficiency anemia in the Makkah region. *Hematology*. 2016;21(6):351-8.
- 140.Rahim F, Saki, N. Age-specific cutoff in discriminating Iron deficiency anemia from beta-thalassemia traits. *IJBC*. 2010;2(4):197-.

### How to Cite This Article:

Jahangiri M, Rahim F, Saki Malehi A, Pezeshki S M S, Ebrahimi M. Differential Diagnosis of Microcytic Anemia, Thalassemia or Iron Deficiency Anemia: A Diagnostic Test Accuracy Meta-Analysis. *Mod Med Lab J*. 2020; 3 (1) :16-29